

## CLAIMS

What is claimed is:

1. An agricultural inoculum suitable for inoculating plant seeds, said inoculum comprising:
  - a fungal antagonist selected from the group consisting of *Trichoderma virens* GL-3 (ATCC 58678) and mutants thereof;
  - a bacterial antagonist selected from the group consisting of *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390) and mutants thereof; and
  - a suitable carrier that is non-phytotoxic, non-bacteriostatic, and non-bacteriocidal.
2. A composition of matter comprising:
  - a plant seed inoculated with a combination comprising a fungal antagonist selected from the group consisting of *Trichoderma virens* GL-3 (ATCC 58678) and mutants thereof and a bacterial antagonist selected from the group consisting of *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390) and mutants thereof;
  - wherein said combination suppresses growth of plant pathogenic fungi.
3. A plant inoculated with a combination comprising:
  - a fungal antagonist selected from the group consisting of *Trichoderma virens* GL-3 (ATCC 58678) and mutants thereof; and
  - a bacterial antagonist selected from the group consisting of *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390) and mutants thereof;

wherein the combination suppresses growth of plant pathogenic fungi and the plant is selected from the group consisting of

corn,

sunflower,

soybean,

field pea, and

wheat.

4. A method of protecting a plant from disease caused by a plant pathogenic fungus comprising:

inoculating seeds from said plant with a combination comprising a fungal antagonist selected from the group consisting of *Trichoderma virens* GL-3 (ATCC 58678) and mutants thereof and a bacterial antagonist selected from the group consisting of *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390) and mutants thereof;

wherein said combination suppresses growth of plant pathogenic fungi.

5. A method of protecting a seed or a plant from disease caused by a plant pathogenic fungus comprising:

inoculating seeds from said plant with a composition comprising a spore-forming fungal antagonist and a spore-forming bacterial antagonist.

6. The method of claim 5 wherein the spore-forming bacterial antagonist is selected from the group *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390) and mutants thereof.

7. A method of protecting a seed or a plant from disease caused by a plant pathogenic fungus comprising:

inoculating seeds from said plant with a composition comprising a fungal antagonist and  
5 a bacterial antagonist;

wherein said combination suppresses growth of plant pathogenic fungi.

8. The method of claim 7 wherein the combination suppresses growth of the plant pathogen  
fungi *Fusarium*, *Phythium*, *Phytophthora* and *Penicillium*.

9. A method of protecting a plant from disease caused by a plant pathogenic fungus comprising:

inoculating seeds from said plant with a composition comprising a fungal antagonist and  
a bacterial antagonist selected from the group consisting of *Bacillus amyloliquefaciens* TJ1000 or  
1BE (ATCC BAA-390) and mutants thereof;

wherein said combination suppresses growth of plant pathogenic fungi.

10. A method for biologically controlling or inhibiting stalk rot or root rot comprising:

coating seeds with an effective amount of a composition comprising *Trichoderma virens*  
GL-3 (ATCC 58678) and *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390).

11. A process for making a composition comprising:

introducing an essentially pure culture of *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390) to a growth medium about eight hours after an essentially pure culture of *Trichoderma virens* GL-3 (ATCC 58678) is introduced to the growth medium; and growing the culture as a competitive culture.

5

12. A process comprising making a composition comprising:

combining an essentially pure culture of *Trichoderma virens* GL-3 (ATCC 58678) with an essentially pure culture of *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390) in a 50:50 mixture; and

applying said composition to a seed at a rate of at least 100,000 spores per seed.

13. A method for protecting plants in a growing medium from damping off and root rot fungal plant disease comprising:

placing in the growing medium in the immediate vicinity of the plant to be protected an effective quantity of the agricultural inoculum of claim 1.

14. A method for protecting plants from fungal plant disease comprising:

adding the composition of claim 2 in an effective quantity to a substrate selected from the group consisting of pelletized calcium sulfate or pelletized lime; and

placing the pellet in the immediate vicinity of the plant to be protected.

15. The method of claim 14 further comprising:

adding another plant growth nutrient to the pellet.

10067165.020102  
10  
15  
20

16. A method for protecting a plant from fungal plant disease comprising:  
adding the agricultural inoculum of claim 1 in an effective quantity to a liquid solution;  
and  
5 applying the liquid solution in the immediate vicinity of the plant.

17. The method of claim 16 further comprising:  
adding an additive to the liquid, said additive being at least one substance selected from  
the group consisting of  
10 a plant nutrient,  
a plant micro-nutrient, and  
a chemical fungicide.

18. A method for biologically controlling a plant disease caused by a plant-colonizing fungus,  
15 the method comprising:  
inoculating a seed of the plant with an effective amount of a microbial inoculant  
comprising a combination of microorganisms having all of the identifying characteristics of  
*Trichoderma virens* Gl-3 (ATCC 58678) and *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC  
BAA-390), said inoculation resulting in the control of said plant disease.

19. The method of claim 18 wherein said inoculation results in the control of more than one  
plant disease.

20. A method comprising:

combining of a spore-forming fungal strain and a spore-forming bacterial strain to produce a product; whereby the combining step enhances ease of use and longevity of the product.

5

21. A method comprising:

applying a *Trichoderma* spp. microorganism and a *Bacillus* spp. microorganism to a wettable powder to produce a combination; and  
applying the combination to a seed.

22. A composition of matter made by combining:

a fungal antagonist selected from the group consisting of *Trichoderma virens* GL-3 (ATCC 58678) and mutants thereof;  
a bacterial antagonist selected from the group consisting of *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390) and mutants thereof; and  
a suitable carrier that is non-phytotoxic, non-bacteriostatic, and non-bacteriocidal.

23. An antagonist for controlling plant pathogens made by combining effective amounts of:

a fungal antagonist selected from the group consisting of *Trichoderma virens* GL-3 (ATCC 58678) and mutants thereof;  
a bacterial antagonist selected from the group consisting of *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390) and mutants thereof; and  
a suitable carrier that is non-phytotoxic, non-bacteriostatic, and non-bacteriocidal.



29. A method for culturing a plant comprising:

applying the antagonist of claim 23 to a seed or to the seedbed of the plant;

planting the seed in the seedbed;

growing the plant to yield a crop; and

harvesting the crop;

wherein said applying step increases the yield of the crop.

30. The method of claim 29 wherein the antagonist is applied to the seed or to the seedbed of a plant selected from the group consisting of

a monocot, and

a dicot.

31. The method of claim 29 wherein the antagonist is applied to the seed or to the seedbed of a plant selected from the group consisting of

a legume plant, and

a non-legume plant.

32. The method of claim 29 wherein the antagonist is applied to the seed or to the seedbed of a plant selected from the group consisting of

corn,

sunflower,

soybean,



field pea, and

wheat.

33. A process comprising:

5 making a composition by combining an essentially pure culture of *Trichoderma virens* GL-3 (ATCC 58678) with an essentially pure culture of *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390) in a mixture; and

10 applying said composition to a seed; wherein said mixture ranges in composition from 10 to 90 percent *Trichoderma virens* GL-3 (ATCC 58678) by culture volume and from 90 to 10 percent *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390) by culture volume.

34. A process comprising:

15 making a composition by combining an essentially pure culture of *Trichoderma virens* GL-3 (ATCC 58678) with a plurality of essentially pure cultures of bacteria in a mixture; and

applying said composition to a seed; wherein said mixture ranges in composition from 10 to 90 percent *Trichoderma virens* GL-3 (ATCC 58678) by culture volume.

35. An antagonist for controlling plant pathogens made by combining effective amounts of:

20 a fungal antagonist selected from the group consisting of a strain of *Trichoderma virens* and mutants thereof;

a bacterial antagonist selected from the group consisting of *Bacillus amyloliquefaciens* TJ1000 or 1BE (ATCC BAA-390) and mutants thereof; and

a suitable carrier that is non-phytotoxic, non-bacteriostatic, and non-bacteriocidal.

